

Generalized Dynamical Spin Chain and 4-Loop Integrability in $\mathcal{N} = 6$ Superconformal Chern-Simons Theory

Dongsu Bak ^a, Hyunsoo Min ^a, Soo-Jong Rey ^{b,c}

a) Physics Department, University of Seoul, Seoul 130-743 KOREA

b) School of Physics and Astronomy, Seoul National University, Seoul 151-747 KOREA

c) Kavli Institute for Theoretical Physics, University of California, Santa Barbara 93106 USA

dsbak@uos.ac.kr, hmin@dirac.uos.ac.kr, sjrey@snu.ac.kr

abstract

We revisit unitary representation of centrally extended $\mathfrak{psu}(2|2)$ excitation superalgebra. We find most generally that ‘pseudo-momentum’, not lattice momentum, diagonalizes spin chain Hamiltonian and leads to generalized dynamic spin chain. All known results point to lattice momentum diagonalization for $\mathcal{N} = 4$ super Yang-Mills theory. Having different interacting structure, we ask if $\mathcal{N} = 6$ superconformal Chern-Simons theory provides an example of pseudo-momentum diagonalization. For $SO(6)$ sector, we study maximal shuffling and next-to-maximal shuffling terms in the dilatation operator and compare them with results expected from $\mathfrak{psu}(2|2)$ superalgebra and integrability. At two loops, we rederive maximal shuffling term (3-site) and find perfect agreement with known results. At four loops, we first find absence of next-to-maximal shuffling term (4-site), in agreement with prediction based on integrability. We next extract maximal shuffling term (5-site), the most relevant term for checking the possibility of pseudo-momentum diagonalization. Curiously, we find that result agrees with integrability prediction based on lattice momentum, as in $\mathcal{N} = 4$ super Yang-Mills theory. Consistency of our results is fully ensured by checks of renormalizability up to six loops.

1 Introduction

The holographic duality between Type IIA string theory on $\text{AdS}_4 \times \mathbb{CP}^3$ and (2+1)-dimensional $\mathcal{N} = 6$ superconformal Chern-Simons theory, discovered by Aharony, Bergman, Jafferis and Maldacena (ABJM) [1], opened a new avenue for exploring the AdS/CFT correspondence [2]. The (2+1)-dimensional superconformal field theory admits Lagrangian formulation, hereafter referred as ABJM theory, so a systematic study between bulk fields and boundary operators is feasible much to the same extent as the holographic duality between Type IIB string theory on $\text{AdS}_5 \times \mathbb{S}^5$ and (3+1)-dimensional $\mathcal{N} = 4$ super Yang-Mills (SYM) theory. The ABJM theory depends on two parameters, level k of the Chern-Simons term and rank N of the gauge group. The planar limit, $N \rightarrow \infty$, of the ABJM theory corresponds to non-interacting limit, $g_s \rightarrow 0$, of the Type IIA string theory. In this limit, the ABJM theory is organized in perturbation theory at weak ‘t Hooft coupling regime $\lambda \equiv (N/k) \ll 1$, while the Type IIA string theory is organized in perturbation theory at strong ‘t Hooft coupling regime $\sqrt{\lambda} \gg 1$.

A very interesting and important aspect of the AdS_4 -ABJM holographic duality is the prospect of ‘integrability’, the feature investigated thoroughly in the AdS_5 –SYM holographic duality [3]-[16]. At strong ‘t Hooft coupling regime, Type IIA string worldsheet dynamics was found integrable at leading order in $1/\sqrt{\lambda}$ [17, 18]. At weak ‘t Hooft coupling regime, $\text{SO}(6)$ sector of the ABJM dilatation operator was found integrable at 2 loops [19, 18]. Extension to full $\text{OSp}(6|4)$ dilatation operator and Bethe ansatz equations were proposed to all orders and numerous consistency checks of the proposal were studied [20] - [27]. It was found that the excitation spectrum in the AdS_4 -ABJM theory is organized by a direct sum of $\mathfrak{psu}(2|2)$ superalgebra. Interestingly, a direct product of the same superalgebra also featured in the excitation spectrum in the AdS_5 -SYM theory [15], [16]. These results all point toward exact ‘quantum integrability’ of the ABJM theory, but a direct analytic proof would be highly desirable.

The purpose of this paper is to study the dilatation operator of the ABJM theory beyond leading order. Our motivations are primarily twofold. First, we wish to understand precise dynamic nature of the alternating spin chain in this theory. In the case of $\mathcal{N} = 4$ super Yang-Mills theory, dilatation operator was described by a novel spin chain whose length changes dynamically [7]. This novelty was in fact a simple consequence of $\mathfrak{psu}(2, 2|4)$ superconformal algebra. For the $\mathfrak{osp}(6|4)$ superconformal algebra of the ABJM theory, we also expect emergence of dynamic spin chain. Magnon excitations in both theories are known to be described by product or sum of *centrally-extended* $\mathfrak{psu}(2|2)$. On the other hand, field contents and their interactions as well as supersymmetry preserved by the spin chain ground-state differ for the two theories. Therefore, precise dynamic nature of the spin chain could also differ each other. Second, we would like to test quantum integrability of the operator contents in this theory. While there were several indications from both weak and strong coupling regime, there was no rigorous proof yet.

We address this problem by comparing spin chain Hamiltonian expected from integrability and weak coupling expansion of it with the dilatation operator computed directly from the ABJM theory at lower but beyond leading orders in perturbation theory.

We begin in section 2 with centrally extended $\mathfrak{psu}(2|2)$ superalgebra, the superalgebra that features both the ABJM and the $\mathcal{N} = 4$ SYM theories. Re-examining the previous study carefully, we find that off-shell magnon excitations, described by unitary representations of the superalgebra, is in general labeled by a ‘pseudo-momentum’ which in general differs from the lattice momentum of the spin chain. Functional relation of the ‘pseudo-momentum’ to the lattice momentum depends on the theory under consideration, on the amount of supersymmetry and order of perturbative interactions. In turn, the ‘pseudo-momentum’ different from the lattice momentum lead to a generalized dynamical spin chain in that the exchange algebra of the magnon excitation is generalized order by order in perturbation theory.

In section 3, focusing on a magnon in $SU(2) \oplus SU(2)$ sector, we compare predictions of quantum integrability to the spectrum of operator contents, equivalently, to general structure of the spin chain Hamiltonian at higher orders in perturbation theory. We focus in the dilatation operator on so-called ‘maximal shuffling’ terms - terms that exchange spins at furthest sites - and ‘next-to-maximal shuffling’ terms - terms that exchange spins one less than furthest sites - at each order, since these terms are the cleanest to compute yet highly nontrivial in structure. We explain diagrammatic origin of these terms up to four loops. At two loops, we rederive maximal shuffling term (3-sites) and find perfect agreement with known results. At four loops, we first find absence of next-to-maximal shuffling term (4-sites), in agreement with integrability. We then extract maximal shuffling term (5-sites).

In section 4, we explain formalism for computing anomalous dimensions from two-point function of single-trace operators in perturbation theory and hence the quantum dilatation operator. We also explain general structure of the dilatation operators and how the maximal shuffling terms of length $2\ell + 1$ and next-to-maximal shuffling terms of order 2ℓ arise diagrammatically at 2ℓ -th order in perturbation theory. In sections 5 and 6, we compute quantum dilatation operator in dimensional regularization. In section 5, we first compute two-loop contribution to two-point function and reproduce the known result of the dilatation operator. We then compute four-loop contribution. Combining both contributions, we extract next-to-maximal shuffling terms at four loops. If quantum integrability holds, result from section 3 indicates these terms should be absent. We indeed find that bosonic and fermionic loop contributions cancel each other – a clear indication that integrability holds. In section 6, we focus on maximal shuffling terms at four loops, where only bosonic diagrams contribute. Again, we find the result agrees with the prediction of integrability based on lattice momentum in section 3. Internal consistency of our results is fully ensured by checks of renormalizability up to six loops.

Along with $\mathcal{N} = 4$ SYM theory, our result escalates a puzzle why the dilatation operators are always diagonalized by lattice momentum eigenstates, not by pseudo-momentum eigenstates. In section 7, we discuss implication of this to other issues pertinent to the ABJM theory. We also discuss various implications of our results. In Appendix A, we present $\mathfrak{psu}(2|2)$ S-matrices in the pseudo-momentum basis. Appendix B is relegated to technical details of precision numerical evaluation of irreducible Feynman diagrams, whose results were used in sections 5 and 6.

2 Off-Shell $\mathfrak{PSU}(2|2)$ and Generalized Dynamic Spin Chain

In the planar limit, we consider a single trace operator of the ABJM scalar fields. In ABJM theory, the scalar fields $Y^I, Y_I^\dagger$ ($I = 1, 2, 3, 4$) transform as $(\mathbf{N}, \bar{\mathbf{N}}; \mathbf{4})$ and $(\bar{\mathbf{N}}, \mathbf{N}; \bar{\mathbf{4}})$ under the $U(N) \times \bar{U}(N)$ gauge group and the $SU(4)$ R-symmetry group. Consider in the planar limit single-trace operators $\mathcal{O}$. By gauge invariance, they form an alternating spin chain, labeled by $SU(4)$ quantum numbers:

$$\mathcal{O}[I_1 I_2 I_3 I_4 I_5 \cdots, I_{2L}] = \text{Tr}(Y^{I_1} Y_{I_2}^\dagger Y^{I_3} Y_{I_4}^\dagger Y^{I_5} \cdots Y_{I_{2L}}^\dagger). \quad (2.1)$$

In the limit $L \rightarrow \infty$, choose the convention that Y^I fields of $\mathbf{4}$ are in the odd sites of the spin chain and $Y_I^\dagger$ of $\bar{\mathbf{4}}$ in the even sites. By group theory, $\mathbf{4}$ at odd sites and $\bar{\mathbf{4}}$ at even sites interact only through δ_j^I .

Rename (Y^1, Y^2, Y^3, Y^4) as $(A_1, A_2, B_2^\dagger, B_1^\dagger)$ where A_a, B_b ($a, b = 1, 2$) transform under the $SU(2)_A$ and the $SU(2)_B$ subgroups of $SU(4)$ [21]. We focus on a restricted set of single-trace operators where A_a and B_b field are placed at odd/ even sites, respectively:

$$\mathcal{O}[a_1 a_2 a_3 a_4 a_5 \cdots] = \text{Tr}(A_{a_1} B_{a_2} A_{a_3} B_{a_4} A_{a_5} \cdots). \quad (2.2)$$

A given operator of this type is mappable to a given spin chain state $|a_1 a_2 a_3 \cdots\rangle$. We shall take the ground state as

$$|O\rangle = |1111\cdots\rangle \quad \leftrightarrow \quad \mathcal{O}[1111\cdots] = \text{Tr}(A_1 B_1 A_1 B_1 A_1 B_1 \cdots). \quad (2.3)$$

For this type of spin chains, there is no interaction between $\mathbf{4}$ and $\bar{\mathbf{4}}$ spins. Hence, the odd-site spin chain and the even-site spin chain behave independently. Then, there will be A -type magnons and B -type magnons propagating independently without interactions between the two types¹.

¹In principle, interaction of double-exchange $\mathbb{P}_{n+1, n+3} \mathbb{P}_{n+2, n+4}$ could arise, where $\mathbb{P}_{ij}$ denotes the permutation operator exchanging spins at the i -th and the j -th sites. In this case, there will be interactions between these two types of magnons.

The magnon excitations are organized by off-shell $\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)$ superalgebra symmetries acting on A - and B -sites, respectively. The $(2|2)$ excitation multiplets are $(A_2, B_2^\dagger | \Psi_1, \Psi_2)$ and $(B_2, A_2^\dagger | \Psi_2^\dagger, \Psi_1^\dagger)$. Therefore, magnons of A - and B -types behave identical. From now on, we shall focus on A -type magnons and denote $(A_2, B_2^\dagger | \Psi_1, \Psi_2)$ as $\Phi = (\phi^1, \phi^2 | \psi^1, \psi^2)$ collectively. One expects a magnon excitation is an eigenstate of lattice momentum of the A -type sites in the alternating spin chain:

$$|p\rangle = \sum_{n=0}^{L-1} e^{inp} |\dots \overbrace{\Phi}^{2n+1} \dots\rangle. \quad (2.4)$$

Here, the ellipses denote spin configurations of the ground state, where A_1 or B_1 in the remaining odd or even sites. Below, we shall suppress denoting these background spins explicit.

The off-shell $\mathfrak{psu}(2|2)$ superalgebra of the excitation symmetry is spanned by the two $\mathfrak{su}(2)$ rotation generators $\mathfrak{R}_b^a, \mathfrak{L}_\beta^\alpha$, the supersymmetry generator $\mathfrak{Q}_a^\alpha$ and the superconformal generator $\mathfrak{S}_\alpha^a$. The off-shell configuration is characterized by $sl(2, \mathbb{R})$ central charges $\mathfrak{C}, \mathfrak{K}, \mathfrak{K}^*$ [15]. Their (anti)commutators are given by [15]

$$\begin{aligned} [\mathfrak{R}_b^a, \mathfrak{J}^c] &= \delta_b^c \mathfrak{J}^a - \frac{1}{2} \delta_b^a \mathfrak{J}^c, \quad [\mathfrak{L}_\beta^\alpha, \mathfrak{J}^\gamma] = \delta_\beta^\gamma \mathfrak{J}^\alpha - \frac{1}{2} \delta_\beta^\alpha \mathfrak{J}^\gamma \\ \{\mathfrak{Q}_a^\alpha, \mathfrak{S}_\beta^b\} &= \delta_a^b \mathfrak{L}_\beta^\alpha + \delta_\beta^b \mathfrak{R}_a^b + \delta_a^b \delta_\beta^\alpha \mathfrak{C} \\ \{\mathfrak{Q}_a^\alpha, \mathfrak{Q}_b^\beta\} &= \varepsilon^{\alpha\beta} \varepsilon_{ab} \mathfrak{K}, \quad \{\mathfrak{S}_\alpha^a, \mathfrak{S}_\beta^b\} = \varepsilon_{\alpha\beta} \varepsilon^{ab} \mathfrak{K}^*. \end{aligned} \quad (2.5)$$

The central charges $\mathfrak{C}$ is related to the energy by $E = \mathfrak{C}$, while $\mathfrak{K}, \mathfrak{K}^*$ introduced at off-shell are related to the momentum. On a state of fundamental representation, the generators act as

$$\mathfrak{R}_b^a |\phi^c\rangle = \delta_b^c |\phi^a\rangle - \frac{1}{2} \delta_b^a |\phi^c\rangle, \quad \mathfrak{L}_\beta^\alpha |\phi^\gamma\rangle = \delta_\beta^\gamma |\phi^\alpha\rangle - \frac{1}{2} \delta_\beta^\alpha |\phi^\gamma\rangle \quad (2.6)$$

and as

$$\begin{aligned} \mathfrak{Q}_a^\alpha |\phi^b\rangle &= a \delta_a^b |\psi^\alpha\rangle \\ \mathfrak{Q}_a^\alpha |\psi^\beta\rangle &= b \varepsilon^{\alpha\beta} \varepsilon_{ab} |\phi^b G(A^+)\rangle \\ \mathfrak{S}_\alpha^a |\phi^b\rangle &= c \varepsilon_{\alpha\beta} \varepsilon^{ab} |\psi^\beta G(A^-)\rangle \\ \mathfrak{S}_\alpha^a |\psi^\beta\rangle &= d \delta_\alpha^\beta |\phi^a\rangle. \end{aligned} \quad (2.7)$$

The function $G(A^\pm)$ is not arbitrary. Note first that A^+ inserts the ground-state spin A_1 into the odd site of the chain while A^- removes one ground-state spin A_1 . Acting on an eigenstate (2.4) of the lattice momentum p , we have

$$G(A^\pm) = A^\pm \quad \text{where} \quad |A^\pm \Phi\rangle = e^{\mp ip} |\Phi A^\pm\rangle. \quad (2.8)$$

This is the defining relation of the new marker function $G(A^\pm)$.

Closure of the superalgebra on fundamental representation leads to the ‘shortening condition’

$$ad - bc = 1. \quad (2.9)$$

One also finds the central charges of total momentum yield

$$\begin{aligned} \mathfrak{K} |\Phi\rangle &= ab |\Phi G(A^+)\rangle \\ \mathfrak{K}^* |\Phi\rangle &= cd |\Phi G(A^-)\rangle \end{aligned} \quad (2.10)$$

and the central charge of total dilatation energy yields

$$\mathfrak{C} |\Phi\rangle = \frac{1}{2}(ad + bc) |\Phi\rangle. \quad (2.11)$$

Quantum mechanically, functional form of $G(A^\pm)$ in (2.7) are subject to radiative corrections. These corrections can be extracted straightforwardly, for example, from explicit derivation of quantum dilatation operator and utilization of the relations (2.10, 2.11) or from operator product expansions between fermionic charges $\mathfrak{Q}, \mathfrak{S}$ and a pair of single-trace operators carrying bosonic and fermionic excitations, respectively. After radiative corrections are taken into account, the full-fledged quantum marker function $G(A^\pm)$ ought to solve the ‘shortening condition’ (2.9). It is an interesting open problem to find a complete representation theoretic classification of $G(A^\pm)$ as an exact, nonperturbative function of λ .

Here, we content ourselves for a particular ansatz of $G(A^\pm)$ motivated by perturbation theory. Introduce a notion of ‘pseudo-momentum’ P by the exchange algebra:

$$P := P(p) \quad \text{where} \quad G(A^\pm)\Phi := e^{\mp iP(p)}\Phi G(A^\pm). \quad (2.12)$$

Physically, we expect that $P(p)$ is an odd function of p and periodic with periodicity 2π . This means that the pseudo-momentum P is expandable in a power series of the lattice momentum p as

$$e^{iP} = G(e^{ip}) = e^{ip} e^{i \sum_{n=1}^{\infty} b_{2n}(\lambda) \sin np}. \quad (2.13)$$

From general structure of the perturbation theory that dictates hopping of an excitation spin over lattice sites, we expect that $b_{2n}(\lambda)$ is further expandable as

$$b_{2n} = \sum_{\ell=n}^{\infty} b_{2\ell, 2n} \lambda^{2\ell}. \quad (2.14)$$

To obey the on-shell conditions $\mathfrak{K} = \mathfrak{K}^* = 0$, one needs to excite two or more spins. Consider a spin chain with n multiple excitation spins that are asymptotically separated and carry ‘pseudo-momentum’ $P_1, \dots, P_n$. The momentum central charges now read

$$\begin{aligned}\mathfrak{K} |\Phi^1 \Phi^2 \Phi^3 \dots \Phi^n\rangle &= K |\Phi^1 \Phi^2 \Phi^3 \dots \Phi^n G(A^+)\rangle \\ \mathfrak{K}^* |\Phi^1 \Phi^2 \Phi^3 \dots \Phi^n\rangle &= K^* |\Phi^1 \Phi^2 \Phi^3 \dots \Phi^n G(A^-)\rangle\end{aligned}\quad (2.15)$$

with

$$K = \sum_{k=1}^n a_k b_k \prod_{m=k+1}^n e^{-iP_m}; \quad K^* = \sum_{k=1}^n c_k d_k \prod_{m=k+1}^n e^{+iP_m}. \quad (2.16)$$

Both should vanish on any on-shell configuration that satisfies

$$\prod_{k=1}^n e^{-iP_k} = 1. \quad (2.17)$$

In terms of the lattice momentum, this in general puts the spin chain to obey a version of twisted boundary condition if closed ². Therefore, one might opt to consider the pseudo-momentum defined only in asymptotic limit. Our point is to stress that the condition at quantum level ought to be the one for the ‘pseudo-momentum’ P , not for the lattice momentum p . As shown in [15], a unique local solution is given by

$$a_k b_k = \alpha(\lambda)(e^{-iP_k} - 1); \quad c_k d_k = \beta(\lambda)(e^{iP_k} - 1). \quad (2.18)$$

Note that α and β are independent of the lattice site k and the lattice momenta p_k ’s – they are parameters *common* to all excitations.

A convenient parametrization of a , b , c and d are in terms of elliptic variables $X^\pm$ and $h(\lambda), \gamma, f$:

$$\begin{aligned}a &= \sqrt{h(\lambda)} \gamma, & b &= -\sqrt{h(\lambda)} \frac{f}{\gamma X^+} (X^+ - X^-) \\ c &= \sqrt{h(\lambda)} \frac{i \gamma}{f X^-}, & d &= -\sqrt{h(\lambda)} \frac{i}{\gamma} (X^+ - X^-).\end{aligned}\quad (2.19)$$

With positive definite $h(\lambda)$, defining relations are

$$\frac{X^+}{X^-} = e^{iP}; \quad (X^+)^* = X^-, \quad (2.20)$$

while the shortening condition (2.9) reads:

$$X^+ + \frac{1}{X^+} - X^- - \frac{1}{X^-} = \frac{i}{h(\lambda)}. \quad (2.21)$$

²We thank N. Beisert for useful correspondences on issues related to this point.

Explicitly,

$$X^\pm = \frac{e^{\pm i\frac{P}{2}}}{4h \sin \frac{P}{2}} \left(1 + \sqrt{1 + 16h^2(\lambda) \sin^2 \frac{P}{2}} \right). \quad (2.22)$$

Unitarity of the representation demands that, modulo a complex phase, $\gamma = \sqrt{-i(X^+ - X^-)}$ and $f = 1$. In terms of these variables, the dilatation energy spectrum reads

$$E = \frac{h}{i}(X^+ - X^-) - \frac{1}{2} = \frac{1}{2} \sqrt{1 + 16h^2(\lambda) \sin^2 \frac{P}{2}}. \quad (2.23)$$

Classically, the ‘pseudo-momentum’ P is reduced to the lattice momentum p , and the elliptic variables $X^\pm$ are reduced to $x^\pm$ where $(x^+/x^-) = e^{ip}$.

As in [15], S-matrices between a pair of magnon excitations is determinable by the $\mathfrak{psu}(2|2)$ superalgebra. For completeness, we tabulate them in appendix A. Following [28], overall scalar phase-factor is also determinable by imposing crossing relations. These S-matrices satisfy unitarity, Yang-Baxter equations and crossing symmetries. From the S-matrices, one can also construct full-fledged Bethe ansatz equations (BAE). We stress that all these conclusions are most transparent when the ‘pseudo-momentum’ P , equivalently, the elliptic variables $X^\pm$ are used instead of the lattice momentum p , equivalently, $x^\pm$.

A remark is in order. Deformation of integrable spin chain was considered in [29], in which the coupling parameter λ is replaced by a set of four deformation parameters. In particular, one of these parameters acts to deform $(x_\pm + \lambda^2/x_\pm) \rightarrow x_\pm + \sum_{n=3} \alpha_n/(x_\pm)^n$ and hence the energy spectrum as well. It was shown that these deformations are possible while retaining global symmetries of the spin chain intact. Such a deformation appears closely related to the map from the lattice momentum to the pseudo-momentum introduced above. It would be extremely interesting to understand possible relation better.

3 Spin Chain Hamiltonian from Integrability

The magnon spectrum expected from the $\mathfrak{psu}(2|2)$ excitation symmetry and the integrability is of the form

$$E = \sqrt{\frac{1}{4} + 4h^2(\lambda) \sin^2 \frac{p}{2}}. \quad (3.1)$$

It was shown that the energy spectrum receives perturbative corrections only from even-loop orders. So, the interpolating function $h^2(\lambda)$ is parametrizable as

$$h^2(\lambda) = \lambda^2 \sum_{\ell=0}^{\infty} h_{2\ell} \lambda^{2\ell} \quad \text{with} \quad h_0 = 1. \quad (3.2)$$

This is the spectrum for the single magnon for each SU(2) sector. Expanding the energy spectrum at weak coupling as a function of lattice momentum p ,

$$\begin{aligned} E &= \frac{1}{2} + 4h^2(\lambda) \sin^2 \frac{p}{2} - 16h^4(\lambda) \sin^4 \frac{p}{2} + \dots \\ &= \left(\frac{1}{2} \right) + \left(4 \sin^2 \frac{p}{2} \right) \lambda^2 + \left(-16 \sin^4 \frac{p}{2} + 4h_2 \sin^2 \frac{p}{2} \right) \lambda^4 + \dots \end{aligned} \quad (3.3)$$

As is well-known from planar perturbation theory, 2ℓ -th order contribution to the energy gives rise to lattice shuffling up to 2ℓ consecutive sites. At 2-loop order, from $e^{\pm ip}$, the maximal lattice shuffling is for 2 sites. At 4-loop order, from $e^{\pm 2ip}$, the maximal lattice shuffling is for 4 sites, etc. We shall refer to these as ‘maximal shuffling’ interactions.

At this stage, as shown in the previous section, one needs to bear in mind of the possibility that dynamic spin chain is diagonalized in the pseudo-momentum P , not in the lattice momentum p . In this case, we need to replace the lattice momentum p in (3.1) by the pseudo-momentum P . As the pseudo-momentum is defined as a function of the lattice momentum in perturbation theory, $P = P(p)$, the energy spectrum is still expandable as double series of the ‘t Hooft coupling and the lattice hopping. Schematically, it takes the form

$$\begin{aligned} E &= \sum_{\ell=0}^{\infty} \lambda^{2\ell} \sum_{n=0}^{\ell} e_{2\ell,2n} \sin^{2n} \frac{p}{2} \\ &= \left(\frac{1}{2} \right) + \left(e_{2,2} \sin^2 \frac{p}{2} \right) \lambda^2 + \left(e_{4,2} \sin^2 \frac{p}{2} + e_{4,4} \sin^4 \frac{p}{2} \right) \lambda^4 + \dots \end{aligned} \quad (3.4)$$

where $e_{0,0} = \frac{1}{2}$ counts the classical scaling dimension and $e_{2\ell,0} = 0$ ($\ell \geq 1$) is fixed by the supersymmetry of the ground-state. The coefficient $e_{2,2} = 4$ was computed previously from explicit computation of quantum dilatation operator at two loop order [19, 18].

The coefficients $e_{2\ell,2n}$ provide a complete information concerning spectra of the quantum dilatation operator, $E(p, \lambda)$. More precisely, from these coefficients, one can determine *both* the generalized marker function $G(e^{ip})$ and the interpolating coupling function $h^2(\lambda)$ as the two functions are mutually independent. Our assertion is based on the observation that, at each order in perturbation theory, the coefficients $e_{2\ell,2n}$ in the spectra are in one-to-one correspondence with the two sets of coefficients $(h_{2\ell}, b_{2\ell,2n})$.

The corresponding integrable Hamiltonian at each order is well-known. Consider the odd-site SU(2) chain. The tree-level part of the Hamiltonian counting the classical scaling dimension is

$$H_0 = \frac{1}{2} \sum_{n=0}^{L-1} \mathbb{I}. \quad (3.5)$$

The 2-loop part of the Hamiltonian is simply given by [19, 18]

$$H_2 = \sum_{n=0}^{L-1} (\mathbb{I} - \mathbb{P}_{a_{2n+1}a_{2n+3}}), \quad (3.6)$$

where $2L$ is the number of the total sites and $\mathbb{P}$ denotes exchange operator. We consider the asymptotic spin chain where L goes to infinity. The 4-loop part of the Hamiltonian can be identified as

$$H_4 = \frac{e_{4,2}}{4} \sum_{n=0}^{L-1} (\mathbb{I} - \mathbb{P}_{a_{2n+1}a_{2n+3}}) + \frac{e_{4,4}}{16} \sum_{n=0}^{L-1} [(\mathbb{P}_{a_{2n+1}a_{2n+5}} - 1) + 4(\mathbb{I} - \mathbb{P}_{a_{2n+1}a_{2n+3}})], \quad (3.7)$$

where we have used the result of the $\mathcal{N} = 4$ super Yang-Mills theory [4, 7, 9]. In order to demonstrate that the energy spectrum (3.4) follows from these Hamiltonian, we use the momentum eigenstate of a single A -magnon excited on the odd-site $SU(2)$ chain:

$$|p\rangle_A = \sum_{n=0}^{L-1} e^{inp} |\dots \overbrace{A_2}^{2n+1} \dots\rangle \quad (3.8)$$

in the background of A_1 s on the ellipses. Likewise, for the even-site $SU(2)$ chain, the momentum eigenstate of a single B -magnon is

$$|p'\rangle_B = \sum_{m=1}^L e^{imp'} |\dots \overbrace{B_2}^{2m} \dots\rangle \quad (3.9)$$

in the background of B_1 s on the ellipses. Therefore, there are four kinds of excitation states:

$$|0\rangle_A \otimes |0\rangle_B, \quad |p\rangle_A \otimes |0\rangle_B, \quad |0\rangle_A \otimes |p'\rangle_B, \quad |p\rangle_A \otimes |p'\rangle_B. \quad (3.10)$$

The second and the fourth states are the momentum p states of the odd-site $SU(2)$ chain. Below, we shall focus on the odd-site chain because the odd-site $SU(2)$ chain and even-site $SU(2)$ chain behave independently for the above quartet of states.

One easily checks that

$$\begin{aligned} H_2 |p\rangle_A &= 4 \sin^2 \frac{p}{2} |p\rangle_A \\ H_4 |p\rangle_A &= \left(e_{4,2} \sin^2 \frac{p}{2} + e_{4,4} \sin^4 \frac{p}{2} \right) |p\rangle_A, \end{aligned} \quad (3.11)$$

etc. In the Hamiltonian (3.7), the maximal shuffling terms $e_{2\ell,2\ell}$ are of particular interest. For example, at 4-loop order, (3.3) indicates that $e_{4,4}$ equals to -16 . In perturbation theory, $e_{4,4}$ in H_4 can be extracted by evaluating anomalous dimensions of a single A -magnon up to 4-loop orders. Since the value $e_{4,4} = -16$ is a consequence of $\mathfrak{psu}(2|2)$ excitation symmetry and the

integrability, explicit computation would amount to a test of quantum integrability of the ABJM theory.

In the following sections, we shall compute the maximal shuffling term $e_{4,4}$ and the next-to-maximal shuffling term $e_{4,3}$ explicitly and compare with prediction of the $\mathfrak{psu}(2|2)$ excitation symmetry and the integrability. For the coefficient $e_{4,4}$ of the maximal shuffling term, one needs to compute the diagram in Fig. 1 and its conjugate. Diagrammatically, there can




Figure 1: *Maximal shuffling: 4-loop interactions involving five sites.*

also arise next-to-maximal shuffling terms. To compute their coefficient $e_{4,3}$, one needs to compute the diagrams in Fig. 2 and Fig. 3. They give rise to dilatation operators of the form $\sum_n (\mathbb{I} - \mathbb{P}_{n+1,n+3})(\mathbb{I} - \mathbb{P}_{n+2,n+4})$. In case one type-A and one type-B magnons are excited simultaneously, this operators represent nontrivial interactions between them. This operator, however, is not diagonalized by the state $|p\rangle_A \otimes |p'\rangle_B$. Absence of such interactions amounts to the statement that $e_{4,3} = 0$.




Figure 2: *Next-to-maximal shuffling: 4-loop interaction diagram involving four sites.*




Figure 3: *Next-to-maximal shuffling: 4-loop interaction diagram involving four sites and fermion loop.*

4 Quantum Dilatation Operator

4.1 Formalism

We use the anomalous dimension matrix that is defined by the two-point correlation function:

$$\langle :O(x): :O(0): \rangle_\epsilon = \frac{C_\epsilon^{2L}}{(x^2)^{\frac{1-\epsilon}{2} \cdot 2L} (x^2)^{\gamma(\epsilon)}} \quad (4.1)$$

where we use the dimensional regularization to control the ultraviolet divergences. This is related to the collections A_n of Feynman diagrams at each order defined by

$$\begin{aligned} \langle :O(x): :O(0): \rangle_\epsilon &= (I_\epsilon)^{2L} e^{-\gamma_\epsilon \ln(x^2 \Lambda^2(\epsilon))} \\ &:= (I_\epsilon)^{2L} \exp[\ln(1 + A_2 \lambda^2 + A_4 \lambda^4 + \dots)] \end{aligned} \quad (4.2)$$

Here, I_ϵ denotes the dimensionally regularized Euclidean scalar propagator in the position space:

$$I_\epsilon = \int \frac{d^{2\omega} p}{(2\pi)^{2\omega}} \frac{1}{p^2} e^{ip \cdot x} = \frac{\Gamma(\omega - 1)}{4\pi^\omega} \frac{1}{(x^2)^{\omega-1}}, \quad (4.3)$$

with $2\omega = 3 - \epsilon$.

From the above, we extract lower loop contributions to the dilatation operator in the ABJM theory³ by

$$\begin{aligned} H_2 &= -\lim_{\epsilon \rightarrow 0} \epsilon A_2 \\ H_4 &= -\lim_{\epsilon \rightarrow 0} 2\epsilon \left(A_4 - \frac{1}{2} A_2^2 \right) \\ H_6 &= -\lim_{\epsilon \rightarrow 0} 3\epsilon \left(A_6 - \frac{1}{2} (A_2 A_4 + A_4 A_2) + \frac{1}{3} A_2^3 \right). \end{aligned} \quad (4.4)$$

Here, the overall factor ℓ in $H_{2\ell}$ arises from extracting the dilatation operator as the charge generating variation of $\ln x^2$ in the ABJM theory. The leading singularity in A_2 starts from the order $O(\epsilon^{-1})$, so the leading term in the Laurent expansion of A_2 contributes to H_2 . For the case of A_4 , the leading singularity in general starts at the $O(\epsilon^{-2})$ order. The coefficient of this leading singularity in $A_4 - \frac{1}{2} A_2^2$ ought to vanish for renormalizability of the theory under consideration. Therefore, the leading singularity starts again from $O(\epsilon^{-1})$ terms. For the case of A_6 , the leading singularity starts at the $O(\epsilon^{-3})$ order. Again, the coefficient of this leading singularity of $O(\epsilon^{-3})$ and of the next-to-leading singularity $O(\epsilon^{-2})$ vanish for renormalizability. This pattern continues to all higher order contributions to the dilatation operator H .

³For perturbative computations of the dilation operator in $\mathcal{N} = 4$ SYM theory in close parallel to the Feynman diagrammatics we take, see [30].

4.2 Maximal Shuffling and Next-To-Maximal Shuffling Interactions

We now focus on the ABJM theory. The operator structure involved with Fig. 1 can be constructed from the operator structure of the two-loop scalar sextet contribution:

$$O_{123}^n = 2\mathbb{I} - \mathbb{K}_{n+1,n+2} - \mathbb{K}_{n+2,n+3} + 2\mathbb{P}_{n+1,n+3}\mathbb{K}_{n+1,n+2} + 2\mathbb{P}_{n+1,n+3}\mathbb{K}_{n+2,n+3} - 4\mathbb{P}_{n+1,n+3} \quad (4.5)$$

where $(\mathbb{K})_{IJ}^{KL} = \delta_{IJ}\delta^{KL}$ refers to contraction operator. The matrix structure of Fig. 1 is then

$$O_{345}^n O_{123}^n = 4\mathbb{I} - 8\mathbb{P}_{n+1,n+3} - 8\mathbb{P}_{n+3,n+5} + 16\mathbb{P}_{n+3,n+5}\mathbb{P}_{n+1,n+3} + \cdots \quad (4.6)$$

where the ellipses denotes the omission of the terms involving any types of contraction between the odd and the even sites. Similarly, there is also a diagram corresponding to the conjugate of Fig. 1. For this, the operator structure takes the form,

$$O_{123}^n O_{345}^n = 4\mathbb{I} - 8\mathbb{P}_{n+1,n+3} - 8\mathbb{P}_{n+3,n+5} + 16\mathbb{P}_{n+1,n+3}\mathbb{P}_{n+3,n+5} + \cdots \quad (4.7)$$

Therefore the full operator structure becomes

$$\begin{aligned} O_{12345}^n &= O_{123}^n O_{345}^n + O_{345}^n O_{123}^n \\ &= 8\mathbb{I} - 16\mathbb{P}_{n+1,n+3} - 16\mathbb{P}_{n+3,n+5} + 16\mathbb{P}_{n+1,n+3}\mathbb{P}_{n+3,n+5} + 16\mathbb{P}_{n+3,n+5}\mathbb{P}_{n+1,n+3} + \cdots \end{aligned} \quad (4.8)$$

If we put only A_a / B_a magnons to the odd- / even-sites of the chain, the above can be rewritten as

$$O_{12345}^n = 16\mathbb{P}_{n+1,n+5} - 8\mathbb{I} + \cdots, \quad (4.9)$$

where we have used the identity

$$\varepsilon_{Ia_1a_3a_5}\varepsilon_{Ib_1b_3b_5} = \mathbb{I} + \delta_{a_1}^{b_3}\delta_{a_3}^{b_5}\delta_{a_5}^{b_1} + \delta_{a_1}^{b_5}\delta_{a_3}^{b_1}\delta_{a_5}^{b_3} - \mathbb{P}_{13} - \mathbb{P}_{35} - \mathbb{P}_{15} = 0. \quad (4.10)$$

The same operator structure arises from $-\frac{1}{2}A_2^2$, the second-order effects of the two-loop contribution to the dilatation operator. Hence, adding these two contributions, one obtains the coefficient $e_{4,4}$.




Figure 4: Remaining 4 loop interaction diagram involving four sites

The operator structure of the diagrams in Fig. 2 involves 4-site interactions and can be extracted as

$$\begin{aligned} O_{1234}^n &= O_{123}^n O_{234}^n + O_{234}^n O_{123}^n \\ &= 32(\mathbb{P}_{n+1,n+3} - \mathbb{I})(\mathbb{P}_{n+2,n+4} - \mathbb{I}) + 8(2\mathbb{P}_{n+1,n+3} + 2\mathbb{P}_{n+3,n+5} - 3\mathbb{I}) + \cdots. \end{aligned} \quad (4.11)$$

Again, the same operator structure arises from $-\frac{1}{2}A_2^2$ contribution to the dilatation operator. Fermions also contribute to the 4-site operators $\mathbb{P}_{n+1,n+3}\mathbb{P}_{n+2,n+4}$. This involves fermion loop as depicted in Fig. 3 and the corresponding matrix structure is given by

$$\bar{O}_{1234}^n = O_{1234}^n + 32\mathbb{I} + \cdots. \quad (4.12)$$

Operators of the form $(\mathbb{P}_{n+1,n+3} - \mathbb{I})(\mathbb{P}_{n+2,n+4} - \mathbb{I})$ contributes to $e_{4,3}$ in H_4 . As said, the $\mathfrak{psu}(2|2)$ excitation symmetry and the integrability predicts that $e_{4,3} = 0$. In the following sections, we shall confirm this explicitly — the contributions of Fig. 2, Fig. 3 and second-order effects of 2-loop contributions all cancel one another.

There are many other diagrams that arise at 4-loops and contribute to lower shuffling operators. Though we shall not evaluate any of these terms in this paper, for completeness of our discussions, we shall list a class of relevant diagrams. The operator structure of Fig. 4 is proportional to

$$Q_{1234}^n = 4\mathbb{P}_{n+1,n+3} + 4\mathbb{P}_{n+2,n+4} - 4\mathbb{I} + \cdots \quad (4.13)$$

Therefore this 4-loop diagram is relevant to the computation of the coefficient $e_{4,2}$.




Figure 5: 4-loop diagram involving 5 sites and only $\mathbb{K}$ operators




Figure 6: 4-loop diagram involving 5 sites and only $\mathbb{K}$ operators




Figure 7: 4-loop diagram involving 4 sites and only $\mathbb{K}$ operators




Figure 8: 4-loop diagram involving 4 sites and fermion loop




Figure 9: 4-loop interaction diagram involving 3 sites




Figure 10: Another 4-loop interaction diagram involving 3 sites

There are many 4-loop diagrams involving only $\mathbb{K}$ operators. Such diagrams involving 5-sites are illustrated in Fig. 5 and Fig. 6 . Further diagrams involving 4-sites are illustrated in Fig. 7 and Fig. 8. The 4-loop diagrams contributing operators of 3-sites are proliferated. In Fig. 9 and Fig. 10, we illustrate two representative diagrams arising from sextet interaction vertices only. They contribute to the coefficient $e_{4,2}$.

5 3-Site and 4-Site Interactions

In this section, we shall first compute contributions to 3-site operators from 2-loop diagrams and 4-site operators from 4-loop diagrams. As discussed in the last section, the second-order effect of the 2-loop diagrams will also contribute to H_4 along with 4-loop diagrams.

5.1 2 loops and 3-site interactions

For the computation of the 4-loop diagrams, we begin with the computation of the 2-loop and the 1-loop diagrams with a momentum flow P . We first compute the 2-loop contribution of Fig. 11 using the dimensional regularization. In this diagram, we suppress symmetry factor first and then reinstate it in the end. We then have

$$L_3(P) = \int \frac{d^{2\omega}p}{(2\pi)^{2\omega}} \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{1}{p^2} \frac{1}{q^2} \frac{1}{(p+q+P)^2} = \frac{(\Gamma(\omega-1))^3 \Gamma(3-2\omega)}{(4\pi)^{2\omega} \Gamma(3\omega-3) (P^2)^{3-2\omega}}. \quad (5.1)$$

To get this result, we have used

$$G(a, b) = \int \frac{d^{2\omega}p}{(2\pi)^{2\omega}} \frac{(4\pi)^\omega (P^2)^{a+b-\omega}}{(p^2)^a ((p+P)^2)^b} = \frac{\Gamma(a+b-\omega) \Gamma(\omega-a) \Gamma(\omega-b)}{\Gamma(a) \Gamma(b) \Gamma(2\omega-a-b)}. \quad (5.2)$$

Similarly, the 1-loop contribution in Fig. 11 is evaluated as

$$L_2(P) = \int \frac{d^{2\omega}p}{(2\pi)^{2\omega}} \frac{1}{p^2} \frac{1}{(p+P)^2} = \frac{(\Gamma(\omega-1))^2 \Gamma(2-\omega)}{(4\pi)^\omega \Gamma(2\omega-2) (P^2)^{2-\omega}}. \quad (5.3)$$



Figure 11: Basic 2-loop and 1-loop diagrams with a momentum flow




Figure 12: 3-loop sextet interaction diagram involving 3 sites

Based on these building blocks, we next compute the diagram of Fig. 12. It is proportional to the operator $\sum O_{123}^n$:

$$K_2 = (I_\epsilon)^3 A_2 = (I_\epsilon)^3 (x^2 \pi)^{3-2\omega} a_2 \sum O_{123}^n, \quad (5.4)$$

where a_2 is given by

$$\begin{aligned} a_2 &= (-(2\pi)^2) \cdot (I_\epsilon)^{-3} (x^2 \pi)^{2\omega-3} \int \frac{d^{2\omega}P}{(2\pi)^{2\omega}} (L_3(P))^2 e^{iP \cdot x} \\ &= -\frac{(2\pi)^2 (\Gamma(\omega-1))^3 (\Gamma(3-2\omega))^2 \Gamma(5\omega-6)}{(4\pi)^3 (\Gamma(3\omega-3))^2 \Gamma(6-4\omega)} \\ &= -\frac{1}{4\epsilon} [1 + \epsilon(1 - \psi(1/2)) + O(\epsilon^2)]. \end{aligned} \quad (5.5)$$

Here, the polygamma function $\psi(1/2)$ takes the value:

$$\psi(1/2) = -\mathbf{C} - 2\ln 2 = -1.963510026\dots \quad (5.6)$$

where $\mathbf{C}$ denotes the Euler's constant. For the Fourier transform, we used the formula,

$$\int \frac{d^{2\omega}p}{(2\pi)^{2\omega}} \frac{1}{(p^2)^\alpha} e^{ip \cdot x} = \frac{\Gamma(\omega - \alpha)}{4^\alpha \pi^\omega \Gamma(\alpha)} \frac{1}{(x^2)^{\omega - \alpha}}. \quad (5.7)$$

We see that the part of the 2-loop Hamiltonian H_2 proportional to $\mathbb{P}_{n+1,n+3}$ is solely coming from this sextet interaction contribution A_2 . Therefore, this part of H_2 becomes

$$H_2^{3s} = -\lim_{\varepsilon \rightarrow 0} \varepsilon a_2 \sum O_{123}^n = \frac{1}{4} \sum O_{123}^n = \sum \left[\frac{1}{2} \mathbb{I} - \mathbb{P}_{n+1,n+3} + \dots \right], \quad (5.8)$$

This reproduces the previous result obtained in Refs. [19, 18].

5.2 4-loops and absence of 4-site interactions

We now turn to the coefficient $e_{4,3}$ of the operator O_{1234}^n . The contributions come from the 4-loop diagram of Fig. 2 and also from $-\frac{1}{2}A_2^2$, the second-order effect of the 2-loop contribution. If the integrability were to hold, contributions to $e_{4,3}$ ought to be zero.

We first compute the 4-loop contribution of Fig. 2. The full contribution including the $(I_\varepsilon)^4$ part takes the form

$$K_4^{4s} = (I_\varepsilon)^4 A_4^{4s} = (I_\varepsilon)^4 (x^2 \pi)^{6-4\varepsilon} b_4 \sum O_{1234}^n. \quad (5.9)$$

To proceed further, we also need the following double integral:

$$J_2 \equiv I_5(1, 3-2\omega, 1, 3-2\omega, 2-\omega), \quad (5.10)$$

where

$$I_5(w_1, w_2, w_3, w_4, w_5) = \int \frac{d^{2\omega}k}{(2\pi)^{2\omega}} \frac{d^{2\omega}l}{(2\pi)^{2\omega}} \frac{(4\pi)^{2\omega} (p^2)^{-2\omega + \sum_{k=1}^5 w_k}}{(k^2)^{w_1} (l^2)^{w_2} ((l-p)^2)^{w_3} ((k-p)^2)^{w_4} ((k-l)^2)^{w_5}} \quad (5.11)$$

The coefficient b_4 can then be computed as

$$b_4 = J_2 \cdot \frac{(\Gamma(\omega-1))^4}{4^4 \pi^2} \left[\frac{\Gamma(3-2\omega)}{\Gamma(3\omega-3)} \right]^2 \cdot \frac{\Gamma(2-\omega)\Gamma(8\omega-10)}{\Gamma(2\omega-2)\Gamma(10-7\omega)}. \quad (5.12)$$

In the Laurent expansion in powers of ε , the integral J_2 starts with the $O(\varepsilon^0)$ order. Therefore, it can be expanded as

$$J_2 = \beta_1 \left[1 + \beta_2 \varepsilon + O(\varepsilon^2) \right]. \quad (5.13)$$

Then b_4 has the expansion

$$b_4 = -\frac{\beta_1}{128\pi\epsilon^2} \left[1 + (\beta_2 - 5 - \psi(1) - 2\psi(1/2))\epsilon + O(\epsilon^2) \right], \quad (5.14)$$

where the polygamma function $\psi(1)$ is given by

$$\psi(1) = -\mathbf{C} = -0.577215\dots \quad (5.15)$$

On the other hand, the corresponding coefficient of O_{1234}^n from $-\frac{1}{2}A_2^2$ is again given by $-\frac{1}{2}a_2^2$. The leading singularity of order $O(\epsilon^{-2})$ contribution from these two ought to cancel each other since the ABJM theory is renormalizable. We shall shortly show that the 4-site contribution of Fig. 3 starts with the order $O(\epsilon^{-1})$. This then the coefficient β_1 to

$$\beta_1 = -4\pi. \quad (5.16)$$

We checked this numerically with high precision in the Appendix B. The corresponding dilatation operator is then given by

$$H_4^b = -\lim_{\epsilon \rightarrow 0} 2\epsilon \left[b_4 - \frac{1}{2}a_2^2 \right] \sum O_{1234}^n = -\frac{1}{16} [\beta_2 - 7 - \psi(1)] \sum O_{1234}^n. \quad (5.17)$$

We found that

$$\beta_2 = 8 + \psi(1) \quad (5.18)$$

by carrying out the integral numerically. Again, see Appendix B. Therefore, we find scalar loop contributions sum up to yield

$$H_4^b = -\frac{1}{16} \sum O_{1234}^n. \quad (5.19)$$

Let us now turn to the contribution from the fermion loop contribution, Fig. 3. There are no more terms that are proportional to O_{1234}^n . Since this diagram is of $O(\epsilon^{-1})$, we may extract its contribution to the dilatation operator directly by computing the diagram in Fig. 13 with zero momentum flow to the amputated external lines. The corresponding Feynman integral is given




Figure 13: 4-loop diagram with amputated external lines

by

$$\begin{aligned}
F_4 &= \int dp dq dk dl \frac{1}{(k-p)^2} \frac{1}{(k-q)^2} \frac{1}{(l-p)^2} \frac{1}{(l-q)^2} \frac{\text{tr } \not{p} \not{q}}{p^2 q^2 k^2} \\
&= \int dp dq dk dl \frac{1}{(k-p)^2} \frac{1}{(k-q)^2} \frac{1}{(l-p)^2} \frac{1}{(l-q)^2} \left(\frac{2}{p^2 k^2} - \frac{(p-q)^2}{p^2 q^2 k^2} \right), \quad (5.20)
\end{aligned}$$

where dp abbreviates for $d^{2\omega}p/(2\pi)^{2\omega}$, etc. Then, reinstating the symmetry factor, the contribution yields

$$A_F = F_4 \cdot \frac{4!}{4!} \cdot (2\pi)^4 \cdot \frac{1}{2} \sum O_{1234}^n \quad (5.21)$$

To compute F_4 , we perform l -integral first, p and q integral next, and finally k -integral using the definition I_5 and $G(a, b)$. One finds

$$F_4 = \frac{1}{(4\pi)^{4\omega}} G(1, 1) G(5 - 3\omega, 1) [2G(2 - \omega, 1) G(4 - 2\omega, 1) - I_5(1, 1, 1, 1, 1 - \omega)]. \quad (5.22)$$

One can evaluate

$$\begin{aligned}
G(1, 1) G(5 - 3\omega, 1) &= \frac{\pi}{\varepsilon} (1 + O(\varepsilon)) \\
2G(2 - \omega, 1) G(4 - 2\omega, 1) &= \frac{4\pi}{\varepsilon} [1 + \varepsilon(3 + \psi(1)) + O(\varepsilon^2)]. \quad (5.23)
\end{aligned}$$

In the Appendix B, we further find numerically that

$$J_3 = I_5(1, 1, 1, 1, 1 - \omega) = \frac{4\pi}{\varepsilon} [1 + \varepsilon(1 + \psi(1)) + O(\varepsilon^2)] \quad (5.24)$$

Hence,

$$A_F = \frac{1}{2\varepsilon} \cdot \frac{1}{32} \sum \bar{O}_{1234}^n \quad (5.25)$$

and the fermion contribution the Hamiltonian H_4 reads

$$H_4^f = 2 \frac{dA_f}{d \ln \Lambda} = \lim_{\varepsilon \rightarrow 0} 2\varepsilon A_F = \frac{1}{16} \sum \bar{O}_{1234}^n. \quad (5.26)$$

Adding the bosonic and the fermionic contributions, the total 4-site contribution becomes

$$H_4^{4s} = H_4^b + H_4^f = \sum (2 \mathbb{I} + \dots), \quad (5.27)$$

and $e_{4,3} = 0$ identically. We see that this conclusion fits exactly with the $\mathfrak{psu}(2|2)$ excitation symmetry and the integrability.

6 5-Site Interactions

We now turn to the 5-site interaction generated by the diagram in Fig. 1.⁴ This computes the coefficient $e_{4,4}$. Below, the double integral (5.11) for some particular choice of $(w_1, \dots, w_5)$ will play a central role. We need to evaluate the integral

$$J_1 \equiv I_5(2 - \omega, 3 - 2\omega, 2 - \omega, 3 - 2\omega, 1). \quad (6.1)$$

The whole contribution of Fig. 1 then takes the form

$$K_4^{5s} = (I_\epsilon)^5 (x^2 \pi)^{6-4\epsilon} A_4^{5s} = (I_\epsilon)^5 (x^2 \pi)^{6-4\epsilon} a_4 \sum O_{12345}^l, \quad (6.2)$$

where

$$a_4 = J_1 \cdot \frac{(\Gamma(\omega - 1))^5}{4^4 \pi^2} \left[\frac{\Gamma(2 - \omega) \Gamma(3 - 2\omega)}{\Gamma(2\omega - 2) \Gamma(3\omega - 3)} \right]^2 \cdot \frac{\Gamma(9\omega - 11)}{\Gamma(11 - 8\omega)}. \quad (6.3)$$

In the Laurent expansion in ϵ , J_1 starts with $O(\epsilon^{-1})$. Therefore, the integral J_1 is expandable as

$$J_1 = \frac{\alpha_1}{\epsilon} \left[1 + \alpha_2 \epsilon + O(\epsilon^2) \right]. \quad (6.4)$$

Then a_4 is expanded as

$$a_4 = -\frac{3\alpha_1 \pi}{64\epsilon^2} \left[1 + (\alpha_2 - 10 - 3\psi(1/2))\epsilon + O(\epsilon^2) \right]. \quad (6.5)$$

On the other hand, $\frac{1}{2}a_2^2$ has the expansion:

$$\frac{1}{2}a_2^2 = \frac{1}{32\epsilon^2} \left[1 + (2 - 2\psi(1/2))\epsilon + O(\epsilon^2) \right]. \quad (6.6)$$

Again, the coefficients of $O(\epsilon^{-2})$ in $a_4 - \frac{1}{2}a_2^2$ must vanish by the renormalizability of the ABJM theory. This determines α_1 as

$$\alpha_1 = -\frac{2}{3\pi}. \quad (6.7)$$

By numerical integration with high precision, we found that α_1 agrees with $-2/(3\pi)$ with high accuracy. Again, see Appendix B. The corresponding contribution to H_4 is then given by

$$H_4^{5s} = -\lim_{\epsilon \rightarrow 0} 2\epsilon \left[a_4 - \frac{1}{2}a_2^2 \right] \sum O_{12345}^n = -\left[\alpha_2 - 12 - \psi(1/2) \right] \left[\sum \mathbb{P}_{n+1, n+5} + \dots \right]. \quad (6.8)$$

⁴We are grateful to Joe Minahan pointing out a potential error to the final conclusion in this section concerning (in)compatibility with lattice momentum.

The coefficient α_2 is computed in the Appendix B as

$$\alpha_2 = 13 + \psi(1/2). \quad (6.9)$$

The H_4^{5s} is thus given by

$$H_4^{5s} = -\sum \mathbb{P}_{n+1, n+5} + \dots. \quad (6.10)$$

Comparing this with the 5-site maximal shuffling term in H_4 , we deduce that

$$e_{4,4} = -16. \quad (6.11)$$

This agrees precisely with the prediction, $e_{4,4} = -16$. This prediction was based on $\mathfrak{psu}(2|2)$ excitation symmetry and the integrability. On the other hand, the prediction was also based on a tacit assumption that the magnon excitation is diagonalized by *lattice* momentum eigenstate. Is it possible that the magnon excitation is actually diagonalized by pseudo-momentum eigenstate? We answer this question in next section.

We close this section with a remark on internal consistency check of the above computations. It is also possible to extract the 6-loop contribution to the Hamiltonian, H_6 in (4.4). There, we first need to make sure that terms of order $O(\varepsilon^{-3})$ and $O(\varepsilon^{-2})$ in $[A_6 - \frac{1}{2}(A_4 A_2 + A_2 A_4) + \frac{1}{3}A_2^3]$ vanish identically by the renormalizability of the ABJM theory. Cancellation of these terms provide a set of stringent consistency test for the computation of a_4 since combinations of a_4 and a_2 (as well as cubic of a_2) ought to cancel against a_6 . The check is straightforward. We confirmed that these singular terms in ε -expansion indeed cancel, ensuring that we computed a_4 correctly. Details of this computation, along with explicit evaluation of H_6 , will be reported in a separate paper.

7 Interpretation and Discussions

7.1 Interpretation

In the last section, explicit computations showed that the 4-loop dilatation operator shows the structure of anticipated maximal and next-to-maximal shuffling term. Moreover, the coefficient of them matched with the value (3.1) dictated by the $\mathfrak{psu}(2|2)$ excitation symmetry and integrability. In this subsection, we compare the result with consideration of pseudo-momentum discussed in section 2. We explain that, as in the $\mathcal{N} = 4$ super Yang-Mills theory, the dilatation operator of the ABJM theory is diagonalized by the lattice momentum eigenstate, not by the pseudo-momentum eigenstate.

To show this, we revisit the energy spectrum (3.1) of a single SU(2) magnon but now assuming that it is an eigenstate of in terms of the pseudo-momentum. Using the relation (2.13)

$$P(p) = p + \sum_{n=1}^{\infty} \sum_{\ell=n}^{\infty} b_{2\ell,2n} \lambda^{2\ell} \sin np \quad (7.1)$$

and the interpolating coupling function (3.2)

$$h^2(\lambda) = \lambda^2 \sum_{\ell=0}^{\infty} h_{2\ell} \lambda^{2\ell} \quad \text{with} \quad h_0 := 1, \quad (7.2)$$

the energy spectrum

$$E = \sqrt{\frac{1}{4} + 4h^2(\lambda) \sin^2 \frac{P(p)}{2}}, \quad (7.3)$$

is now expanded in terms of the lattice momentum as

$$E = \frac{1}{2} + \left[4 \sin^2 \frac{p}{2}\right] \lambda^2 + \left[-(16 + 8b_{2,2}) \sin^4 \frac{p}{2} + (4h_2 + 8b_{2,2}) \sin^2 \frac{p}{2}\right] \lambda^4 + \dots \quad (7.4)$$

This falls into the pattern we conjectured for the most general spectrum (3.4):

$$\begin{aligned} E &= \sum_{\ell=0}^{\infty} \lambda^{2\ell} \sum_{n=0}^{\ell} e_{2\ell,2n} \sin^{2n} \frac{p}{2} \\ &= \left(\frac{1}{2}\right) + \left(e_{2,2} \sin^2 \frac{p}{2}\right) \lambda^2 + \left(e_{4,2} \sin^2 \frac{p}{2} + e_{4,4} \sin^4 \frac{p}{2}\right) \lambda^4 + \dots \end{aligned} \quad (7.5)$$

with

$$e_{2,2} = 4, \quad e_{4,2} = 4h_2 + 8b_{2,2}, \quad e_{4,4} = -16 - 8b_{2,2}, \dots$$

One notes that shift from lattice momentum to pseudo-momentum affects the coefficient of the maximal shuffling term. From the last relation ⁵, one learns that our computation (6.11) amounts $b_{2,2} = 0$, viz. lattice momentum eigenstate diagonalizes the dilatation operator.

It is easy to see that the identification works uniquely at each order in perturbation theory. The coefficients $e_{2\ell,2n}$ of conjectured terms in the energy spectrum (3.4) is in one-to-one correspondence with the two sets of coefficients $(h_{2\ell}, b_{2\ell,2n})$. Since the latter two originate from totally different functions, the interpolating coupling function $h^2(\lambda)$ and the generalized marker function $G(z)$, we see that the spectrum $E(p, \lambda)$ determines these two functions uniquely.

⁵The first relation corresponds to the 2-loop results [19] [18]. The second relation carries useful information regarding the interpolating function $h(\lambda)$ to the first nontrivial order in the weak coupling perturbation theory.

7.2 Discussions

In this paper, we studied realization of $\mathfrak{psu}(2|2)$ excitation symmetry and potential integrability in the ABJM theory. The theory is very different from the $\mathcal{N} = 4$ super Yang-Mills theory, only sharing these two aspects. Given this, we raised the possibility that eigenstates (not just eigenvalues) diagonalizing the dilatation operator are not the lattice momentum basis but the more general pseudo-momentum basis whose functional form $P(p)$ is subject to perturbative corrections. Our result shows the contrary: the ABJM theory is essentially the same as the $\mathcal{N} = 4$ SYM theory, at least in these aspects.

Our result confirms that the ABJM theory is quantum integrable up to 4-loop order in weak coupling perturbation theory. We focused on the $SU(2) \times SU(2)$ inside $SO(6)$ scalar sector and on maximal shuffling operators in the Hamiltonian, but we expect the integrability extends to the full $OSp(6|4)$ and to all operators straightforwardly, as in [20]. We believe our result adds further evidence for exact quantum integrability of the ABJM theory. Below, we discuss implications of our results and issues that deserve further investigation.

- Our consideration in section 2 indicates that notion of pseudo-momentum basis is ubiquitous for the off-shell $\mathfrak{psu}(2|2)$ superalgebra [15]. This superalgebra features both the $\mathcal{N} = 4$ SYM theory and the ABJM theory, but since contents and details of the two theories are very different, precise form of the pseudo-momentum $P(p)$ would differ for the two theories. Nevertheless, result of this paper indicates that the dilatation operators of both theories are diagonalized by lattice momentum eigenbasis. The fact that change to pseudo-momentum basis did not actually take place in both theories poses a puzzle and additional hidden structure yet to be understood better. A possible checkpoint would be a direct computation of the off-shell central charges $\mathfrak{R}, \mathfrak{R}^*$ in weak coupling perturbation theory. By comparing the result with the local solution (2.18), one may be able to understand why all $b_{2\ell, 2n}$ vanish in these two theories.
- Our result also bears an implication to the interpolating function $h^2(\lambda)$. The leading perturbative correction h_2 in (7.2) is extractable, for instance, from 3-site shuffling term $e_{4,2}$ in the 4-loop spectrum⁶. From (7.6), one sees that $e_{4,2}$ is directly related to h_2 if $b_{2,2}$ vanishes and the dilatation operator is diagonalized by the lattice momentum eigenstates.
- The notion of pseudo-momentum also calls for revisiting the interpretation of the giant magnon at strong 't Hooft coupling limit for both AdS_5 [32] and AdS_4 [27] cases.. In the light-cone gauge of the *Lorentzian* string worldsheet, it is always possible to

⁶The computation is currently under progress by J. Minahan, O. Ohlson Sax and C. Sieg (private communications). See also [31].

reparametrize each chiral worldsheet coordinates separately. This implies that there can in general be an arbitrariness in identifying the worldsheet momentum conjugate to the chiral worldsheet coordinate with the angular separation of the string configuration in the giant magnon.

- Extension of our result to the parity-violating $\mathcal{N} = 6$ superconformal Chern-Simons theory with gauge group $U(M) \times U(N)$ is straightforward. Two-loop Hamiltonian was constructed in [33] and was found integrable. Our result in this paper indicates that the integrability persists up to four loops: one only needs to replace λ^2 to the geometric mean of $\lambda^2(M)$ and $\lambda^2(N)$.
- By introducing a variety of D-branes wrapping supersymmetric cycles in $\mathbb{CP}^3$, one can construct open spin chains. For AdS_5 / SYM_4 correspondence, several open spin chains including those ending on giant gravitons were found quantum integrable [34]. The $\mathfrak{psu}(2|2)$ superalgebra or appropriate subalgebra continued to act as the excitation symmetry. Similarly, we expect that integrable open spin chains exist in $ABJ(M)$ theories by adding D-branes and flavors [35]. With concrete realization and identification of excitation symmetries therein, we expect that one can learn more not only about the quantum integrability but also aspects of the pseudo-momentum in the presence of boundaries.
- Extension to higher loop orders would be highly desirable. It will teach us not only the magnon energy spectrum and quantum integrability but also other pertinent issues including wrapping interactions, structural (dis)similarity of the dilatation operator with the $\mathcal{N} = 4$ SYM theory etc.

We are currently investigating these issues and intend to report further results in separate publications.

Acknowledgement

We are grateful to Niklas Beisert, David J. Gross, Joe Minahan, Didina Serban and Matthias Staudacher for very helpful discussions and correspondences. SJR thanks warm hospitality of the Kavli Institute for Theoretical Physics during this work. This work was supported in part by Grants from the Korea National Science Foundation R01-2008-000-10656-0, SRC-CQeST-R11-2005-021, 2008-313-C00175, 2005-084-C00003, 2009-008-0372, EU-FP Marie Curie Research & Training Networks HPRN-CT-2006-035863 (2009-06318) and the U.S. National Science Foundation under Grant No. PHY05-51164 at KITP.

A $\mathfrak{psu}(2|2)$ S-matrices with Pseudo-Momentum

In this section we record the $SU(2|2)$ invariant S matrix. Only change is to replace $x^\pm$ of S matrix in [15] by $X^\pm$ which have more complicated momentum dependence. The S matrix reads⁷

$$\begin{aligned}
s_{12}|\phi_1^a\phi_2^b\rangle &= A_{12}|\phi_2^{\{a}\phi_1^{b\}}\rangle + B_{12}|\phi_2^{[a}\phi_1^{b]}\rangle + \frac{1}{2}C_{12}\epsilon^{ab}\epsilon_{\alpha\beta}|\psi_2^\alpha\psi_1^\beta\bar{G}(A^-)\rangle \\
s_{12}|\psi_1^\alpha\psi_2^\beta\rangle &= D_{12}|\psi_2^{\{\alpha}\psi_1^{\beta\}}\rangle + E_{12}|\psi_2^{[\alpha}\psi_1^{\beta]}\rangle + \frac{1}{2}F_{12}\epsilon^{\alpha\beta}\epsilon_{ab}|\phi_2^a\phi_1^bG(A^+)\rangle \\
s_{12}|\phi_1^a\psi_2^\beta\rangle &= G_{12}|\psi_2^\beta\phi_1^a\rangle + H_{12}|\phi_2^a\psi_1^\beta\rangle \\
s_{12}|\psi_1^\alpha\phi_2^b\rangle &= K_{12}|\psi_2^\alpha\phi_1^b\rangle + L_{12}|\phi_2^b\psi_1^\alpha\rangle
\end{aligned} \tag{A.1}$$

where

$$\begin{aligned}
A_{12} &= S_{12}^0 \frac{X_2^+ - X_1^-}{X_2^- - X_1^+} \\
B_{12} &= S_{12}^0 \frac{X_2^+ - X_1^-}{X_2^- - X_1^+} \left(1 - 2 \frac{1 - 1/(X_2^- X_1^+)}{1 - 1/(X_2^- X_1^-)} \frac{X_2^+ - X_1^+}{X_2^+ - X_1^-} \right) \\
C_{12} &= S_{12}^0 \frac{2\gamma_1\gamma_2/(X_2^- X_1^-)}{1 - 1/(X_2^- X_1^-)} \frac{X_2^+ - X_1^+}{X_2^- - X_1^+} \\
D_{12} &= -S_{12}^0 \\
E_{12} &= -S_{12}^0 \left(1 - 2 \frac{1 - 1/(X_2^+ X_1^-)}{1 - 1/(X_2^+ X_1^+)} \frac{X_2^- - X_1^-}{X_2^- - X_1^+} \right) \\
F_{12} &= S_{12}^0 \frac{2\gamma_1\gamma_2/(X_2^+ X_1^+)}{1 - 1/(X_2^+ X_1^+)} \frac{X_2^- - X_1^-}{X_2^- - X_1^+} \\
G_{12} &= S_{12}^0 \frac{X_2^+ - X_1^+}{X_2^- - X_1^+} \\
H_{12} &= S_{12}^0 \frac{\gamma_1}{\gamma_2} \frac{X_2^+ - X_2^-}{X_2^- - X_1^+} \\
K_{12} &= S_{12}^0 \frac{\gamma_2}{\gamma_1} \frac{X_1^+ - X_1^-}{X_2^- - X_1^+} \\
L_{12} &= S_{12}^0 \frac{X_2^- - X_1^-}{X_2^- - X_1^+}.
\end{aligned} \tag{A.2}$$

The phase S_{12}^0 satisfying $S_{12}^0 S_{21}^0 = 1$ will be specified below. The above is for the odd chain S matrices. For the even chain one can have $2|2$ excitations transforming under the $su(2|2)$

⁷More precisely one needs to include the correction in Ref.[14].

symmetry. The S matrix for the $N = 6$ Chern-Simons theory is consisting of

$$\begin{aligned} S^{oo} &= S^{ee} = S_{12}^0 \hat{S} \\ S^{oe} &= S^{eo} = \tilde{S}_{12}^0 \hat{S}. \end{aligned} \quad (\text{A.3})$$

where $\hat{S}$ denotes the matrix part of the above S matrix without the phase S_{12}^0 . One also finds the crossing symmetric phases by

$$\begin{aligned} S_{12}^0 &= \frac{1 - 1/(X_2^+ X_1^-)}{1 - 1/(X_2^- X_1^+)} \sigma(X_1, X_2) \\ \tilde{S}_{12}^0 &= \frac{X_2^- - X_1^+}{X_2^+ - X_1^-} \sigma(X_1, X_2) \end{aligned} \quad (\text{A.4})$$

where $\sigma(X_1, X_2)$ is the dressing phase found in [13].

The full S matrices above satisfy the unitarity condition

$$S_{12}^f S_{21}^f = I \quad (\text{A.5})$$

and the Yang-Baxter equation

$$S_{12}^f S_{13}^f S_{23}^f = S_{23}^f S_{13}^f S_{12}^f. \quad (\text{A.6})$$

We note that these structures relies only on local exchange algebras, independent global boundary conditions such as (2.17)

B Numerical Evaluation of the integrals J_1 , J_2 and J_3

In this appendix, we evaluate the two-loop integrals $I_5(w_1, w_2, w_3, w_4, w_5)$ in (5.11). It is difficult to find an analytically closed form of them. However there exist several packages with which we can numerically evaluate them. It turns out that the Mathematica packages, MB[37, 38] and AMBRE[39] are useful. Since these packages utilize a method based on the Mellin-Barnes representation of integrals, we cast the integrals into the form:

$$\begin{aligned} I_5(w_1, w_2, w_3, w_4, w_5) &= \int \frac{dz_1}{2\pi i} \int \frac{dz_2}{2\pi i} \frac{\Gamma(-z_1) \Gamma(\omega - w_{25} - z_1) \Gamma(\omega - w_1 + z_1)}{\Gamma(w_1 - z_1)} \\ &\quad - \times \frac{\Gamma(-z_2) \Gamma(\omega - w_{35} - z_2) \Gamma(\omega - w_4 + z_2)}{\Gamma(w_4 - z_2)} \\ &\quad \times \frac{\Gamma(-\omega + w_{14} - z_1 - z_2) \Gamma(-\omega + w_{235} + z_1 + z_2) \Gamma(w_5 + z_1 + z_2)}{\Gamma(w_2) \Gamma(w_3) \Gamma(w_5) \Gamma(w_{235}) \Gamma(2\omega - w_{235}) \Gamma(2\omega - w_{14} + z_1 + z_2)}, \end{aligned} \quad (\text{B.1})$$

where w_{235} denotes $w_2 + w_3 + w_5$. Note that this corresponds to a three dimensional version of the expression (25) in [36]. After modifying the package AMBRE to be applicable to three

dimensions, a direct application of these packages produces the numerical value of the desired function

$$\begin{aligned} J_1 &= I_5(2 - \omega, 3 - 2\omega, 2 - \omega, 3 - 2\omega, 1) = \frac{\alpha_1}{\epsilon} [1 + \alpha_2 \epsilon + O(\epsilon^2)] \\ &= -\frac{0.212207\dots}{\epsilon} [1 + (11.0365\dots)\epsilon + O(\epsilon^2)] \end{aligned} \quad (\text{B.2})$$

From this, one may easily read $\alpha_1 = -\frac{2}{3\pi}$ and $\alpha_2 = 13 + \psi(\frac{1}{2})$. For the integral J_2 , we have found the following numerical value

$$\begin{aligned} J_2 &= I_5(1, 3 - 2\omega, 1, 3 - 2\omega, 2 - \omega) = \beta_1 [1 + \beta_2 \epsilon + O(\epsilon^2)] \\ &= -(12.5664\dots) [1 + (7.42278\dots)\epsilon + O(\epsilon^2)]. \end{aligned} \quad (\text{B.3})$$

This is consistent with the expressions $\beta_1 = -4\pi$ and $\beta_2 = 8 + \psi(1)$. The last integral is evaluated as

$$\begin{aligned} J_3 &= I_5(1, 1, 1, 1, 1 - \omega) = \frac{12.5664\dots}{\epsilon} [1 + (0.422784\dots)\epsilon + O(\epsilon^2)] \\ &= \frac{4\pi}{\epsilon} [1 + (1 + \psi(1))\epsilon + O(\epsilon^2)]. \end{aligned} \quad (\text{B.4})$$

References

- [1] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, *N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, JHEP **0810** (2008) 091 [arXiv:0806.1218 [hep-th]].
- [2] J. M. Maldacena, *The large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113] [arXiv:hep-th/9711200].
- [3] J. A. Minahan and K. Zarembo, *The Bethe-ansatz for N = 4 super Yang-Mills*, JHEP **0303** (2003) 013 [arXiv:hep-th/0212208].
- [4] N. Beisert, C. Kristjansen and M. Staudacher, *The dilatation operator of N = 4 super Yang-Mills theory*, Nucl. Phys. B **664** (2003) 131 [arXiv:hep-th/0303060].
- [5] I. Bena, J. Polchinski and R. Roiban, *Hidden symmetries of the AdS(5) × S⁵ superstring*, Phys. Rev. D **69** (2004) 046002 [arXiv:hep-th/0305116].
- [6] N. Beisert and M. Staudacher, *The N = 4 SYM integrable super spin chain*, Nucl. Phys. B **670** (2003) 439 [arXiv:hep-th/0307042].

- [7] N. Beisert, *The $su(2|3)$ dynamic spin chain*, Nucl. Phys. B **682**, 487 (2004) [arXiv:hep-th/0310252].
- [8] D. Serban and M. Staudacher, *Planar $N = 4$ gauge theory and the Inozemtsev long range spin chain*, JHEP **0406** (2004) 001 [arXiv:hep-th/0401057].
- [9] B. Eden, C. Jarczak and E. Sokatchev, *A three-loop test of the dilatation operator in $N = 4$ SYM*, Nucl. Phys. B **712**, 157 (2005) [arXiv:hep-th/0409009].
- [10] M. Staudacher, *The factorized S -matrix of CFT/AdS*, JHEP **0505** (2005) 054 [arXiv:hep-th/0412188].
- [11] N. Beisert and M. Staudacher, *Long-range $PSU(2,2|4)$ Bethe ansatz for gauge theory and strings*, Nucl. Phys. B **727** (2005) 1 [arXiv:hep-th/0504190].
- [12] A. Rej, D. Serban and M. Staudacher, *Planar $N = 4$ gauge theory and the Hubbard model*, JHEP **0603** (2006) 018 [arXiv:hep-th/0512077].
- [13] N. Beisert, R. Hernandez and E. Lopez, *A crossing-symmetric phase for $AdS(5) \times S^5$ strings*, JHEP **0611**, 070 (2006) [arXiv:hep-th/0609044]; N. Beisert, B. Eden and M. Staudacher, *Transcendentality and crossing*, J. Stat. Mech. **0701**, P021 (2007) [arXiv:hep-th/0610251].
- [14] G. Arutyunov, S. Frolov and M. Zamaklar, *The Zamolodchikov-Faddeev algebra for $AdS(5) \times S^5$ superstring*, JHEP **0704**, 002 (2007) [arXiv:hep-th/0612229].
- [15] N. Beisert, *The $su(2|2)$ dynamic S -matrix*, Adv. Theor. Math. Phys. **12**, 945 (2008) [arXiv:hep-th/0511082].
- [16] N. Beisert, *The Analytic Bethe Ansatz for a Chain with Centrally Extended $su(2|2)$ Symmetry*, J. Stat. Mech. **0701** (2007) P017 [arXiv:nlin/0610017].
- [17] G. Arutyunov and S. Frolov, *Superstrings on $AdS_4 \times CP^3$ as a Coset Sigma-model*, JHEP **0809**, 129 (2008) [arXiv:0806.4940 [hep-th]]
 B. j. Stefanski, *Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$* , Nucl. Phys. B **808**, 80 (2009) [arXiv:0806.4948 [hep-th]]
 N. Gromov and P. Vieira, *The AdS_4/CFT_3 algebraic curve*, JHEP **0902** (2009) 040 [arXiv:0807.0437 [hep-th]]
 J. Gomis, D. Sorokin and L. Wulff, *The complete $AdS_4 \times CP^3$ superspace for the type IIA superstring and D-branes*, JHEP **0903** (2009) 015 [arXiv:0811.1566 [hep-th]].

- [18] D. Bak and S. J. Rey, *Integrable Spin Chain in Superconformal Chern-Simons Theory*, JHEP **0810**, 053 (2008) [arXiv:0807.2063 [hep-th]].
- [19] J. A. Minahan and K. Zarembo, *The Bethe ansatz for superconformal Chern-Simons*, JHEP **0809**, 040 (2008) [arXiv:0806.3951 [hep-th]].
- [20] N. Gromov and P. Vieira, *The all loop AdS_4/CFT_3 Bethe ansatz*, JHEP **0901** (2009) 016 [arXiv:0807.0777 [hep-th]]:
C. Ahn and R. I. Nepomechie, *$N=6$ super Chern-Simons theory S-matrix and all-loop Bethe ansatz equations*, JHEP **0809**, 010 (2008) [arXiv:0807.1924 [hep-th]]:
P. Sundin, *The $AdS_4 \times CP^3$ string and its Bethe equations in the near plane wave limit*, JHEP **0902**, 046 (2009) [arXiv:0811.2775 [hep-th]].
- [21] D. Gaiotto, S. Giombi and X. Yin, *Spin Chains in $N=6$ Superconformal Chern-Simons-Matter Theory*, arXiv:0806.4589 [hep-th]
G. Grignani, T. Harmark and M. Orselli, *The $SU(2) \times SU(2)$ sector in the string dual of $N=6$ superconformal Chern-Simons theory*, arXiv:0806.4959 [hep-th]
- [22] G. Papathanasiou and M. Spradlin, *The Morphology of $N=6$ Chern-Simons Theory*, arXiv:0903.2548 [hep-th].
- [23] B. I. Zwiebel, *Two-loop Integrability of Planar $N=6$ Superconformal Chern-Simons Theory*, arXiv:0901.0411 [hep-th]
J. A. Minahan, W. Schulgin and K. Zarembo, *Two loop integrability for Chern-Simons theories with $N=6$ supersymmetry*, arXiv:0901.1142 [hep-th]
C. Ahn and R. I. Nepomechie, *Two-loop test of the $N=6$ Chern-Simons theory S-matrix*, arXiv:0901.3334 [hep-th].
- [24] T. Nishioka and T. Takayanagi, *On Type IIA Penrose Limit and $N=6$ Chern-Simons Theories*, JHEP **0808** (2008) 001 [arXiv:0806.3391 [hep-th]]
D. Astolfi, V. G. M. Puletti, G. Grignani, T. Harmark and M. Orselli, *Finite-size corrections in the $SU(2) \times SU(2)$ sector of type IIA string theory on $AdS_4 \times CP^3$* , arXiv:0807.1527 [hep-th]
- [25] T. McLoughlin and R. Roiban, *Spinning strings at one-loop in $AdS_4 \times P^3$* , JHEP **0812** (2008) 101 [arXiv:0807.3965 [hep-th]]
L. F. Alday, G. Arutyunov and D. Bykov, *Semiclassical Quantization of Spinning Strings in $AdS_4 \times CP^3$* , JHEP **0811** (2008) 089 [arXiv:0807.4400 [hep-th]]
C. Krishnan, *AdS_4/CFT_3 at One Loop*, JHEP **0809** (2008) 092 [arXiv:0807.4561 [hep-th]]

- T. McLoughlin, R. Roiban and A. A. Tseytlin, *Quantum spinning strings in $AdS_4 \times CP^3$: testing the Bethe Ansatz proposal*, JHEP **0811** (2008) 069 [arXiv:0809.4038 [hep-th]].
- [26] K. Zarembo, *Worldsheet spectrum in $AdS(4)/CFT(3)$ correspondence*, arXiv:0903.1747 [hep-th].
- [27] G. Grignani, T. Harmark, M. Orselli and G. W. Semenoff, *Finite size Giant Magnons in the string dual of $N=6$ superconformal Chern-Simons theory*, JHEP **0812**, 008 (2008) [arXiv:0807.0205 [hep-th]]
 I. Shenderovich, *Giant magnons in AdS_4/CFT_3 : dispersion, quantization and finite-size corrections*, arXiv:0807.2861 [hep-th]
 R. Suzuki, *Giant Magnons on CP^3 by Dressing Method*, arXiv:0902.3368 [hep-th]
 M. C. Abbott, I. Aniceto and O. O. Sax, *Dyonic Giant Magnons in CP^3 : Strings and Curves at Finite J* , arXiv:0903.3365 [hep-th].
- [28] R. A. Janik, *The $AdS(5) \times S^5$ superstring worldsheet S -matrix and crossing symmetry*, Phys. Rev. D **73** (2006) 086006 [arXiv:hep-th/0603038].
- [29] N. Beisert and T. Klose, J. Stat. Mech. **0607** (2006) P006 [arXiv:hep-th/0510124]
 T. Bargheer, N. Beisert and F. Loebbert, J. Stat. Mech. **0811** (2008) L11001 [arXiv:0807.5081 [hep-th]]
 T. Bargheer, N. Beisert and F. Loebbert, arXiv:0902.0956 [hep-th].
- [30] D. J. Gross, A. Mikhailov and R. Roiban, *Operators with large R charge in $N = 4$ Yang-Mills theory*, Annals Phys. **301** (2002) 31 [arXiv:hep-th/0205066].
- [31] J. Minahan, talk at Integrability in Gauge and String Theory, Aug. 11 - 15, 2008, Utrecht, Netherland (<http://www.science.uu.nl/IGST08/video/Minahan.html>).
- [32] D. M. Hofman and J. M. Maldacena, *Giant magnons*, J. Phys. A **39** (2006) 13095 [arXiv:hep-th/0604135]
 G. Arutyunov, S. Frolov and M. Zamaklar, *Finite-size effects from giant magnons*, Nucl. Phys. B **778**, 1 (2007) [arXiv:hep-th/0606126].
- [33] D. Bak, D. Gang and S. J. Rey, *Integrable Spin Chain of Superconformal $U(M) \times U(N)$ Chern-Simons Theory*, JHEP **0810**, 038 (2008) [arXiv:0808.0170 [hep-th]].
- [34] D. M. Hofman and J. M. Maldacena, *Reflecting magnons*, JHEP **0711** (2007) 063 [arXiv:0708.2272 [hep-th]]
 H. Y. Chen and D. H. Correa, *Comments on the Boundary Scattering Phase*, arXiv:0712.1361 [hep-th]

- C. Ahn, D. Bak and S. J. Rey, *Reflecting Magnon Bound States*, JHEP **0804**, 050 (2008) [arXiv:0712.4144 [hep-th]]
- D. H. Correa and C. A. S. Young, *Reflecting magnons from D7 and D5 branes*, J. Phys. A **41** (2008) 455401 [arXiv:0808.0452 [hep-th]]
- D. Bak, *Zero Modes for the Boundary Giant Magnons*, Phys. Lett. B **672**, 284 (2009) [arXiv:0812.2645 [hep-th]].
- [35] D. Gaiotto and D. L. Jafferis, *Notes on adding D6 branes wrapping RP^3 in $AdS_4 \times CP^3$* , arXiv:0903.2175 [hep-th]
- Y. Hikida, W. Li and T. Takayanagi, *ABJM with Flavors and FQHE*, arXiv:0903.2194 [hep-th]
- S. Hohenegger and I. Kirsch, *A note on the holography of Chern-Simons matter theories with flavour*, arXiv:0903.1730 [hep-th].
- [36] I. Bierenbaum and S. Weinzierl, *The massless two-loop two-point function*, Eur. Phys. J. C **32**, 67 (2003) [arXiv:hep-ph/0308311].
- [37] M. Czakon, *Automatized analytic continuation of Mellin-Barnes integrals*, Comput. Phys. Commun. **175**, 559 (2006) [arXiv:hep-ph/0511200].
- [38] A. V. Smirnov and V. A. Smirnov, *On the Resolution of Singularities of Multiple Mellin-Barnes Integrals*, arXiv:0901.0386 [hep-ph].
- [39] J. Gluza, K. Kajda and T. Riemann, *AMBRE - a Mathematica package for the construction of Mellin-Barnes representations for Feynman integrals*, Comput. Phys. Commun. **177**, 879 (2007) [arXiv:0704.2423 [hep-ph]].